

Lecture Notes for

Logic and Computability

Course Number: IND04033UF

by

Bettina Könighofer

Bernd Grabner

Belinda Uhl

Contact

Bettina Könighofer

Institute for Applied Information Processing and Communications (IAIK)

Graz University of Technology, Austria

bettina.koenighofer@iaik.tugraz.at

March, 2021



Graz University of Technology

Table of Contents

7	Natural Deduction for Predicate Logic	1
7.1	Natural Deduction Rules	1
7.1.1	Proof Rules for Universal Quantification	1
7.1.2	Proof Rules for Existential Quantification	3
7.1.3	Proof Rules for Equality	5
7.1.4	Quantifier Equivalences	6
7.1.5	Counterexamples	7
7.2	Self-Evaluation	9
7.2.1	Examples	9
7.2.2	Solutions	10
7.2.3	Self-Evaluation	14

7

Natural Deduction for Predicate Logic

In this chapter, we will discuss the natural deduction calculus for predicate logic. The proofs are formed by using the rules that we already discussed for propositional logic and additionally we add new rules for the quantifiers and equality. As in the natural deduction calculus for propositional logic, we will discuss *introduction* and *elimination* rules for the *quantifiers* and the *equality* predicate.

7.1 Natural Deduction Rules

7.1.1 Proof Rules for Universal Quantification

The ‘ \forall -Elimination’ Rule

First we discuss the rule for eliminating \forall :

$$\frac{\forall x \phi}{\phi [t/x]} \forall e$$

The rule says that if $\forall x \phi$ is true, then we could replace the x in ϕ with any term t (with the side condition for substitution that t has to be free for x in ϕ) and conclude that $\phi [t/x]$ is also true. Recall that $\phi [t/x]$ is obtained by replacing all free occurrences of x in ϕ by t . Since ϕ is assumed to be true for all x , then ϕ should also be true for any term t .

The ‘ \forall -Introduction’ Rule

Now, let us take a look at the rule for introducing a \forall :

$$\frac{\begin{array}{|l} x_0 \\ \vdots \\ \phi [x_0/x] \end{array} \quad x_0 \text{ fresh}}{\forall x \phi} \quad \forall i$$

This rule is a bit more complicated. Just like we have seen in propositional logic, it is composed of a proof box. The box used in this rule defines the scope of the *freshly introduced variable* x_0 ; i.e., the box limits the usage of x_0 to this box only. *The proof-rule tells us, that if we are starting with a **fresh variable** x_0 and we are able to prove some formula $\phi [x_0/x]$, we are able to derive $\forall x \phi$.*

Note, that x_0 has to be a fresh variable, that has never been used outside the box. As it is a fresh variable, we can consider it as an arbitrary term, of which we did not assume anything before. Thus any other term would also work in its place (e.g., x_1, y_2, t, \dots). The rule says that if you can prove ϕ about an x_0 that is free and therefore is not special in any way, then you could prove it for any x whatsoever, hence the conclusion $\forall x \phi$.

Example. Give the proof for the following sequent:

$$\forall x (P(x) \rightarrow Q(x)), \forall x P(x) \vdash \forall x Q(x).$$

Solution.

1.	$\forall x (P(x) \rightarrow Q(x))$	prem.
2.	$\forall x P(x)$	prem.
3.	$x_0 \quad P(x_0) \rightarrow Q(x_0)$	$\forall e \ 1$
4.	$P(x_0)$	$\forall e \ 2$
5.	$Q(x_0)$	$\rightarrow e \ 3,4$
6.	$\forall x Q(x)$	$\forall i \ 3-5$

The structure of this proof is guided by the fact that the conclusion is a \forall formula, therefore the application of the $\forall i$ rule is needed. So we set up the box controlling the scope of x_0 , and we need to prove $Q(x_0)$ inside the box in order to be able to conclude $\forall x Q(x)$ outside of the box. Using $\forall e$, we get the two instances of the premises $P(x_0)$ and $P(x_0) \rightarrow Q(x_0)$ used to prove $Q(x_0)$.

Example. Give the proof for the following sequent:

$$\forall x (\neg P(x) \rightarrow Q(x)), \neg Q(t) \vdash P(t)$$

Solution.

1. $\forall x (\neg P(x) \rightarrow Q(x))$ prem.
2. $\neg Q(t)$ prem.
3. $\neg P(t) \rightarrow Q(t)$ $\forall e$ 1
4. $\neg\neg P(t)$ MT 3,2
5. $P(t)$ $\neg\neg e$ 4

Note that if you apply the $\forall e$ rule, you can use for the substitution any term t (free for x in ϕ) which is helpful in your current proof.

7.1.2 Proof Rules for Existential Quantification

The ‘ \exists -Introduction’ Rule

The $\exists i$ rule is simply:

$$\frac{\phi [t/x]}{\exists x \phi} \exists i$$

The rule tells us, that if $\phi [t/x]$ is true, we can conclude $\exists x \phi$. As $\exists x$ only asks for ϕ to be true for some term t (naturally, we impose the side condition that t be free for x in ϕ).

The ‘ \exists -Elimination’ Rule

The rule for eliminating an \exists relates to the already known $\forall e$. The $\exists e$ rule is defined as follows:

$$\frac{\begin{array}{c} \boxed{\begin{array}{c} x_0 \\ \phi [x_0/x] \text{ ass.} \\ \vdots \\ \chi \end{array}} \quad x_0 \text{ fresh} \\ \exists x \phi \quad \chi \quad \exists e \end{array}}$$

Just like when eliminating \forall , we need to make a case analysis. As $\exists x \phi$ is assumed to be true, we know that ϕ is true for at least one ‘value’ of x , but we do not know for which value. So we do a case analysis over all those possible values by using a **fresh variable** x_0 representing all of those values. So if $\phi [x_0/x]$ allows us to prove some χ (that does not contain x_0), χ can be deduced outside of the box. The box is controlling two things: the scope of x_0 and also the scope of the assumption $\phi[x_0/x]$.

Example. Give the proof for the following sequent:

$$\forall x (P(x) \rightarrow Q(x)), \exists x P(x) \vdash \exists x Q(x)$$

Solution.

1.	$\forall x (P(x) \rightarrow Q(x))$	prem.
2.	$\exists x P(x)$	prem.
3.	$x_0 P(x_0)$	ass.
4.	$P(x_0) \rightarrow Q(x_0)$	$\forall e$ 1
5.	$Q(x_0)$	$\rightarrow e$ 4,3
6.	$\exists x Q(x)$	$\exists i$ 5
7.	$\exists x Q(x)$	$\exists e$ 2,3-6

The motivation for introducing the box in line 3 of this proof is the existential quantifier in the premise $\exists x P(x)$ which has to be eliminated. In line 4 we eliminate the \forall from line 1. Now, we can extract $Q(x_0)$ using line 4 and line 3. In line 6 we introduce an \exists and substitute the x_0 again with an x .

As the formula in line 6 does not contain x_0 any more, we now may close the box in accordance to our $\exists e$ rule. To conclude our $\exists e$, which we started with the box at line 3, in line 7 we need to rewrite the same formula as in line 6.

Example. Consider the following proof and analyse the error made in this proof:

1.	$\forall x (P(x) \rightarrow Q(x))$	prem.
2.	$\exists x P(x)$	prem.
3.	$x_0 P(x_0)$	ass.
4.	$P(x_0) \rightarrow Q(x_0)$	$\forall e$ 1
5.	$Q(x_0)$	$\rightarrow e$ 4,3
6.	$Q(x_0)$	$\exists e$ 2,3-5
7.	$\exists x Q(x)$	$\exists i$ 6

Solution. Line 6 allows the fresh variable x_0 to escape the scope of the box which declares it. This is illegal. Therefore, the $\exists i$ rule has to be applied already inside of the box like in the proof above.

Boxes may also be nested within each other. But we need to be careful, on where our scopes begin and where they end. To understand the concept of multiple boxes, we take a look at another interesting example.

Example. Give the proof for the following sequent:

$$\exists x P(x), \forall x \forall y (P(x) \rightarrow Q(y)) \vdash \forall y Q(y)$$

Solution.

1.	$\exists x P(x)$	prem.
2.	$\forall x \forall y (P(x) \rightarrow Q(y))$	prem.
3.	y_0	
4.	$x_0 P(x_0)$	ass.
5.	$\forall y (P(x_0) \rightarrow Q(y))$	$\forall e$ 2
6.	$P(x_0) \rightarrow Q(y_0)$	$\forall e$ 5
7.	$Q(y_0)$	$\rightarrow e$ 6,4
8.	$Q(y_0)$	$\exists e$ 1,4-7
9.	$\forall y Q(y)$	$\forall i$ 3-8

In this example, the first premise is an \exists formula, which requires an $\exists e$ to be of any use. The conclusion is an \forall formula, which requires the application of the $\forall i$ rule. Therefore, this proof has two boxes. The outer box from 3-8 is for introducing \forall , whereas the inner box from 4-7 is for eliminating the \exists from line 1. We need to declare for both boxes fresh variables. To keep it simple, we will substitute y_0 for y for the outer box and x_0 for x for the inner box. Note again, that it is important to not use x_0 and y_0 outside of these boxes.

7.1.3 Proof Rules for Equality

The ‘= -Introduction’ Rule

It holds that any term t has to be equal to itself. This is expressed by the introduction rule for equality:

$$\frac{}{t=t} =i$$

The rule is not very useful on its own, but in combination with the $=e$ rule it is (similarly to the combination of LEM and $\forall e$).

The ‘= -Elimination’ Rule

This rule allows us, to substitute equals for equals. For example, suppose that $y * (w + 2)$ equals $(y * w) + (y * 2)$ then it is the case that $z \geq y * (w + 2)$ implies $z \geq (y * w) + (y * 2)$. The elimination rule for equality is defined as follows:

$$\frac{t_1 = t_2 \quad \phi [t_1/x]}{\phi [t_2/x]} =e$$

Note that t_1 and t_2 have to be free for x in ϕ , whenever we want to apply the rule $=e$.

Example. Give the proof for the following sequent:

$$(a + 1) = (1 + a), (a + 1 > 1) \rightarrow (a + 1 > 0) \vdash (1 + a > 1) \rightarrow (1 + a > 0)$$

Solution.

1. $(a + 1) = (1 + a)$ prem.
2. $(a + 1 > 1) \rightarrow (a + 1 > 0)$ prem.
3. $(1 + a > 1) \rightarrow (1 + a > 0)$ = e 1,2

In this proof we have $(a + 1)$ is t_1 , $(1 + a)$ is t_2 , and $(x > 1) \rightarrow (x > 0)$ is ϕ . The first premise tells us that t_1 is equal to t_2 . If we now take a look at the second premise, we can see that our formula ϕ is already in the form of $\phi [t_1/x]$. Given this knowledge, we are able to use the two premises to substitute t_1 with t_2 to form our conclusion in line 3.

Example. Give the proof for the following sequent:

$$\forall x (x = f(y)) \vdash \forall y (y = f(x))$$

Solution.

1. $\forall x (x = f(y))$ prem.
2.

y_0	$y_0 = f(y)$	$\forall e$ 1
3.	$f(x) = f(y)$	$\forall e$ 1
4.	$y_0 = f(x)$	$= e$ 2,3
5. $\forall y (y = f(x))$ $\forall i$ 2-4

Since the conclusion is a \forall formula, we apply the $\forall i$ rule and open a box with a fresh variable y_0 . The trick in this example is to apply the $\forall e$ rule twice on the first premise: once by substituting y_0 for x and once using $f(x)$ for x . Next we apply the $= e$ rule. In this case, t_1 is $f(x)$, t_2 is $f(y)$ and ϕ is $y_0 = k$. As $\phi [f(x)/k]$ is already stated in line 2, we are able to conclude line 4 by substituting k with $f(y)$.

7.1.4 Quantifier Equivalences

To train to perform your own natural deduction proofs, you can consider the creating proofs for the most commonly used quantifier equivalences. The proofs are interesting, because most of them involve several quantifications over more than just one variable and your proofs will have nested boxes.

For example, you can proof the following equivalences by creating a proof for each direction:

$$\begin{aligned}
 \neg \forall x \phi &\equiv \exists x \neg \phi \\
 \neg \exists x \phi &\equiv \forall x \neg \phi \\
 \neg \forall x \neg \phi &\equiv \exists x \phi \\
 \neg \exists x \neg \phi &\equiv \forall x \phi
 \end{aligned}$$

Example. Proof the following quantifier equivalence:

$$\neg\exists x P(x) \equiv \forall x \neg P(x)$$

Solution. For both directions, we create a proof:

$$\forall x \neg P(x) \vdash \neg\exists x P(x)$$

$$\neg\exists x P(x) \vdash \forall x \neg P(x)$$

1. $\forall x \neg P(x)$ prem.

2. $\exists x P(x)$ ass.

3. $t \quad P(t)$ ass.

4. $\neg P(t)$ $\forall e$ 1

5. \perp $\neg e$ 3,4

6. \perp $\exists e$ 2,3-5

7. $\neg\exists x P(x)$ $\neg i$ 2-6

1. $\neg\exists x P(x)$ prem.

2. t

3. $P(t)$ ass.

4. $\exists x P(x)$ $\exists i$ 3

5. \perp $\neg e$ 1,4

6. $\neg P(t)$ $\neg i$ 3-5

7. $\forall x \neg P(x)$ $\forall i$ 2-6

7.1.5 Counterexamples

If a sequent is not valid, there is no natural deduction proof for such a sequent. In such cases, we construct a counter-example that proves the sequent to be invalid. As discussed in the chapters addressing propositional logic, a *counter-example* is a model, that satisfies all the premises but does not satisfy the conclusion.

Example. Show that the following sequent is invalid by constructing a counter-example for it:

$$\exists x (P(x) \rightarrow S) \vdash \exists x P(x) \rightarrow S$$

Solution. We define the following model \mathcal{M} that serves as a counterexample:

- $\mathcal{A} = \{a, b\}$
- $P^{\mathcal{M}} = \{a\}$
- $S^{\mathcal{M}} = \perp$

Note, that the \exists operator binds strongest. Therefore the scope of $\exists x$ is only $P(x)$ in the conclusion.

To show, that \mathcal{M} , we first show that \mathcal{M} violates the conclusion, and second that it satisfies the premise.

Showing that $\mathcal{M} \not\models \exists x P(x) \rightarrow S$: First we draw the syntax tree as shown in Figure 7.1. Then we evaluate the nodes. To satisfy the $\exists x$ node in our formula, we need to find at least one value for x that $P(x)$ true. When substituting $[a/x]$, we get that $P^{\mathcal{M}}(a) = \text{true}$, which also makes the $\exists x$ node true. The predicate S always evaluates to false. Therefore, the implication results in a \perp , thus making the conclusion false.




Figure 7.1: Syntax tree of the conclusion. Red nodes are *false*, green nodes are *true*. Dotted and dashed nodes are subtrees with substituted variables.

Showing that $\mathcal{M} \models \exists x (P(x) \rightarrow S)$: Again, we draw the syntax tree as shown in Figure 7.2 and evaluate the nodes. In order for the $\exists x$ node to become true, we need to find a value for x that makes the implication node true. Again we first try to substitute $[a/x]$, which results in a true $P(x)$, but in a false implication. If we, however, substitute $[b/x]$, $P(x)$ evaluates to false and thus making the implication true. Thus also our $\exists x$ is true and therefore also the whole premise.




Figure 7.2: Syntax tree of the premise. Red nodes are *false*, green nodes are *true*. Dotted and dashed nodes are subtrees with substituted variables.

We now have shown, that our given model \mathcal{M} satisfies the premise, but not the conclusion. Therefore \mathcal{M} is a counterexample. If we would not have been able to find a counterexample, the sequence must have been true.

7.2 Self-Evaluation

7.2.1 Examples

For each of the following sequences, either provide a natural deduction proof, or a counter example, that proves the sequent invalid. For proofs, clearly indicate which rule, and what assumptions, premises or intermediate results you are using in each step. Also clearly indicate the scope of any boxes you use. For counterexamples, give a complete model. Show that the model satisfies the premise(s) of the sequent in question, but does not satisfy the respective conclusion.

Example 1. $\forall x (P(x) \wedge Q(x)) \vdash \forall x (P(x) \vee Q(x))$

Example 2. $\forall x (P(x) \rightarrow Q(x)) \vdash \exists y (P(y) \rightarrow Q(y))$

Example 3. $\forall x \forall y P(x, y) \vdash \forall a \forall b P(a, b)$

Example 4. $\exists x (P(x) \rightarrow Q(x)) \vdash \forall x (P(x) \rightarrow Q(x))$

Example 5. $\forall x (P(x) \wedge Q(x)) \vdash \forall x P(x) \wedge \forall x Q(x)$

Example 6. $\exists x P(x) \vdash \neg \forall x \neg P(x)$

Example 7. $P(b) \vdash \forall x (x = b \rightarrow P(x))$

Example 8. $\exists x (P(x) \rightarrow Q(y)), \exists x P(x) \vdash Q(y)$

Example 9. $\forall x P(a, x, x), \forall x \forall y \forall z (P(x, y, z) \rightarrow P(f(x), y, f(z)))$
 $\vdash P((f(a), a, f(a)))$

Example 10. $\exists y \forall x (s(x) = t(y)) \vdash \forall x \exists y (s(x) = t(y))$

Example 11. $\forall x \exists y (s(x) = t(y)) \vdash \exists y \forall x (s(x) = t(y))$

Example 12. $\forall x \forall y (\neg P(x, y) \rightarrow \neg(y = x)) \vdash \forall x P(x, x) \vee \forall y P(y, y)$

Example 13. $\neg \forall x (P(x) \wedge Q(x) \wedge R(y)) \vdash \exists x \neg(P(x) \wedge Q(x) \wedge R(y))$

Example 14. $\exists x \neg(P(x) \wedge Q(x) \wedge R(y)) \vdash \neg \forall x (P(x) \wedge Q(x) \wedge R(y))$

7.2.2 Solutions

Solution 1. $\forall x (P(x) \wedge Q(x)) \vdash \forall x (P(x) \vee Q(x))$

1. $\forall x (P(x) \wedge Q(x))$ prem.
2. $P(x_0) \wedge Q(x_0)$ $\forall e$ 1
3. $P(x_0)$ $\wedge e_1$ 2
4. $P(x_0) \vee Q(x_0)$ $\vee i_1$ 3
5. $\exists x (P(x) \vee Q(x))$ $\exists i$ 4

Solution 2. $\forall x (P(x) \rightarrow Q(x)) \vdash \exists y (P(y) \rightarrow Q(y))$

1. $\forall x (P(x) \rightarrow Q(x))$ prem.
2. $P(t) \rightarrow Q(t)$ $\forall e$ 1
3. $\exists y (P(y) \rightarrow Q(y))$ $\exists i$ 2

Solution 3. $\forall x \forall y P(x, y) \vdash \forall a \forall b P(a, b)$

1. $\forall x \forall y P(x, y)$ prem.
2. $x_0 \quad \forall y P(x_0, y)$ $\forall e$ 1
3. $y_0 \quad P(x_0, y_0)$ $\forall e$ 2
4. $\forall b P(x_0, b)$ $\forall i$ 3
5. $\forall a \forall b P(a, b)$ $\forall i$ 4

Solution 4. $\exists x (P(x) \rightarrow Q(x)) \vdash \forall x (P(x) \rightarrow Q(x))$

Model M :

- $A = \{a, b\}$
- $P^M = \{a\}$
- $Q^M = \perp$

M satisfies premise, but not conclusion.

M is a counterexample.

Solution 5. $\forall x (P(x) \wedge Q(x)) \vdash \forall x P(x) \wedge \forall x Q(x)$

1.	$\forall x (P(x) \wedge Q(x))$	prem.
2.	$x_0 P(x_0) \wedge Q(x_0)$	$\forall e$ 1
3.	$P(x_0)$	$\wedge e_1$ 2
4.	$\forall x P(x)$	$\forall i$ 2-3
5.	$y_0 P(y_0) \wedge Q(y_0)$	$\forall e$ 1
6.	$Q(y_0)$	$\wedge e_2$ 5
7.	$\forall x Q(x)$	$\forall i$ 5-6
8.	$\forall x P(x) \wedge \forall x Q(x)$	$\wedge i$ 4,7

Solution 6. $\exists x P(x) \vdash \neg \forall x \neg P(x)$

1.	$\exists x P(x)$	prem.
2.	$\forall x \neg P(x)$	ass.
3.	$x_0 P(x_0)$	ass.
4.	$\neg P(x_0)$	\forall 2
5.	\perp	$\neg e$ 3,4
6.	\perp	$\exists e$ 1,4-5
7.	$\neg \forall x \neg P(x)$	$\neg i$ 2-6

Solution 7. $P(b) \vdash \forall x (x = b \rightarrow P(x))$

1.	$P(b)$	prem.
2.	x_0	
3.	$x_0 = b$	ass.
4.	$P(x_0)$	$= e$ 3,1
5.	$(x_0 = b) \rightarrow P(x_0)$	$\rightarrow i$ 3-4
6.	$\forall x (x = b \rightarrow P(x))$	$\forall i$ 2-5

Solution 8. $\exists x (P(x) \rightarrow Q(y)), \exists x P(x) \vdash Q(y)$

Model M :

- $A = \{a, b\}$
- $P^M = \{b\}$
- $Q^M = \perp$

M satisfies premises, but not conclusion.

M is a counterexample.

Solution 9. $\forall x P(a, x, x), \forall x \forall y \forall z (P(x, y, z) \rightarrow P(f(x), y, f(z)))$
 $\vdash P((f(a), a, f(a)))$

1. $\forall x P(a, x, x)$ prem.
2. $\forall x \forall y \forall z (P(x, y, z) \rightarrow P(f(x), y, f(z)))$ prem.
3. $P(a, a, a)$ $\forall e$ 1
4. $\forall y \forall z (P(a, y, z) \rightarrow P(f(a), y, f(z)))$ $\forall e$ 2
5. $\forall z (P(a, a, z) \rightarrow P(f(a), a, f(z)))$ $\forall e$ 4
6. $P(a, a, a) \rightarrow P(f(a), a, f(a))$ $\forall e$ 5
7. $P((f(a), a, f(a)))$ $\rightarrow e$ 6,3

Solution 10. $\exists y \forall x (s(x) = t(y)) \vdash \forall x \exists y (s(x) = t(y))$

1. $\exists y \forall x (s(x) = t(y))$ prem.
2.

y_0	$\forall x (s(x) = t(y_0))$	ass.
-------	-----------------------------	------
3.

x_0	$s(x_0) = t(y_0)$	$\forall e$ 2
-------	-------------------	---------------
4.

$\exists y (s(x_0) = t(y))$	$\exists i$ 3
-----------------------------	---------------
5.

$\forall x \exists y (s(x) = t(y))$	$\forall i$ 3-4
-------------------------------------	-----------------
6. $\forall x \exists y (s(x) = t(y))$ $\exists e$ 1,2-5

Solution 11. $\forall x \exists y (s(x) = t(y)) \vdash \exists y \forall x (s(x) = t(y))$

Model M :

- $A = \{1, 2\}$
- $s(x) = \begin{cases} 1 & \text{if } x = 1 \\ 2 & \text{if } x = 2 \end{cases}$
- $t(y) = \begin{cases} 1 & \text{if } y = 1 \\ 2 & \text{if } y = 2 \end{cases}$

M satisfies premises, but not conclusion.

M is a counterexample.

Solution 12. $\forall x \forall y (\neg P(x, y) \rightarrow \neg(y = x)) \vdash \forall x P(x, x) \vee \forall y P(y, y)$

1.	$\forall x \forall y (\neg P(x, y) \rightarrow \neg(y = x))$	prem.
2.	$t \quad \forall y (\neg P(t, y) \rightarrow \neg(y = t))$	$\forall e$ 1
3.	$\neg P(t, t) \rightarrow \neg(t = t)$	$\forall e$ 2
4.	$t = t$	= i
5.	$\neg \neg t = t$	$\neg \neg i$ 4
6.	$\neg \neg P(t, t)$	MT 3,5
7.	$P(t, t)$	$\neg \neg e$ 6
8.	$\forall x P(x, x)$	$\forall i$ 2-7
9.	$\forall x P(x, x) \vee \forall y P(y, y)$	$\forall i_1$ 8

Solution 13. $\neg \forall x (P(x) \wedge Q(x) \wedge R(y)) \vdash \exists x \neg (P(x) \wedge Q(x) \wedge R(y))$

1.	$\neg \forall x (P(x) \wedge Q(x) \wedge R(y))$	prem.
2.	$P(t) \wedge Q(t) \wedge R(y)$	ass.
3.	$\forall x (P(x) \wedge Q(x) \wedge R(y))$	$\forall i$ 2
4.	\perp	$\neg e$ 1,3
5.	$\neg (P(t) \wedge Q(t) \wedge R(y))$	$\neg i$ 2-4
6.	$\exists x \neg (P(x) \wedge Q(x) \wedge R(y))$	$\exists i$ 5

Solution 14. $\exists x \neg (P(x) \wedge Q(x) \wedge R(y)) \vdash \neg \forall x (P(x) \wedge Q(x) \wedge R(y))$

1.	$\exists x \neg (P(x) \wedge Q(x) \wedge R(y))$	prem.
2.	$\forall x (P(x) \wedge Q(x) \wedge R(y))$	ass.
3.	$t \quad \neg (P(t) \wedge Q(t) \wedge R(y))$	ass.
4.	$P(t) \wedge Q(t) \wedge R(y)$	$\forall e$ 2
5.	\perp	$\neg e$ 3,4
6.	\perp	$\exists e$ 1,3-5
7.	$\neg \forall x (P(x) \wedge Q(x) \wedge R(y))$	$\neg i$ 2-6

7.2.3 Self-Evaluation

These examples will be solved online during class.

Example 1. $\neg\forall x\neg P(x) \vdash \exists xP(x)$

Example 2. $f(x, y) = x, g(x, y) = y \vdash f(f(x, y), g(x, y)) = x$

Example 3. $\forall x(A(x) \rightarrow \neg B(x)) \wedge \neg\neg\forall xC(x) \vdash$
 $(\exists x\neg A(x) \vee \exists x\neg B(x)) \wedge \forall xC(x)$

Example 4. $P(y) \rightarrow \forall xQ(x), \exists x\neg Q(x) \vdash \exists x\neg P(x)$

Example 5. $\forall xP(x) \wedge \forall xQ(x) \vdash \forall x(P(x) \wedge Q(x))$

Example 6. $\exists xP(x) \vee \exists xQ(x) \vdash \exists x(P(x) \vee Q(x))$

Declaration of Sources

Chapter 7 was based on the following book.

Michael Huth, Mark Dermot Ryan: *Logic in Computer Science: Modelling and Reasoning about Systems*. June 2004. Cambridge University Press. ISBN:978-0-521-54310-1